

Summary of the Kolmogorov-Obukhov (1941) theory: overview.

Summary of the Kolmogorov-Obukhov (1941) theory: overview.

In the last three posts we have summarised various aspects of the Kolmogorov-Obukhov (1941) theory. When considering this theory, the following things need to be borne in mind.

[a] Whether we are working in x -space or k -space matters. See my posts of 8 April and 15 April 2021 for a concise general discussion.

[b] In x -space the equation of motion (NSE) simply presents us with the problem of an insoluble, nonlinear partial differential equation.

[c] In k -space the NSE presents a problem in statistical physics and in itself tells us much about the transfer and dissipation of turbulent kinetic energy.

[d] The Karman-Howarth equation is a local energy balance that holds for any particular value of the distance r between two measuring points.

[e] There is no energy flux between different values of r ; or, alternatively, through scale.

[f] The energy flux $\Pi(k)$ is derived from the Lin equation (i.e. in wavenumber space) and cannot be applied in x -space.

[g] The maximum value of the energy flux, $\Pi_{\max} = \varepsilon_T$ (say), is a number, not a function, and can be used (like the dissipation ε) in both k -space and x -space.

[h] It also matters whether the isotropic turbulence we are considering is stationary or decaying in time.

[g] If the turbulence is decaying in time, then K41B relies on Kolmogorov's hypothesis of *local stationarity*. It has been pointed out in a previous post (Part 2 of the present series)

that this cannot be the case by virtue of restriction to a range of scales nor in the limit of infinite Reynolds number [1]. See also the supplemental material for [2].

[h] In k -space this is not a problem and the $k^{-5/3}$ spectrum can still be expected [1], as of course is found in practice.

[i] If the turbulence is stationary, then K41B is exact for a range of wavenumbers for sufficiently large Reynolds numbers. The extent of this inertial range increases with increasing Reynolds numbers.

I have not said anything in this series about the concept of intermittency corrections or anomalous exponents. This topic has been dealt with in various blogs and soon will be again.

[1] W. D. McComb and R. B. Fairhurst. The dimensionless dissipation rate and the Kolmogorov (1941) hypothesis of local stationarity in freely decaying isotropic turbulence. J. Math. Phys., 59:073103, 2018.

[2] W. David McComb, Arjun Berera, Matthew Salewski, and Sam R. Yoffe. Taylor's (1935) dissipation surrogate reinterpreted. Phys. Fluids, 22:61704, 2010.